




THEOREM OF THE DAY



The Change of Variables Theorem Let A be a region in \mathbb{R}^2 expressed in coordinates x and y . Suppose that region B in \mathbb{R}^2 , expressed in coordinates u and v , may be mapped onto A via a 1 – 1 transformation T specified by continuously differentiable functions $x = p(u, v)$ and $y = q(u, v)$. Then for a continuous function f on A ,

$$\iint_A f dx dy = \iint_B f(p(u, v), q(u, v)) |J_T(u, v)| du dv,$$

where $J_T(u, v) = \det \begin{pmatrix} \partial p / \partial u & \partial p / \partial v \\ \partial q / \partial u & \partial q / \partial v \end{pmatrix}$ is the Jacobian of T .



Do you know what a mathematician is? A mathematician is one to whom *that* is as obvious as that twice two makes four is to you!



Lord Kelvin's integral on the blackboard offers a famous example of how today's theorem can dramatically simplify a problem. Let $I = \int_{-\infty}^{\infty} e^{-x^2} dx$. Then $I^2 = (\int_{-\infty}^{\infty} e^{-x^2} dx)(\int_{-\infty}^{\infty} e^{-y^2} dy) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy$. We propose to change variable by expressing the xy plane in polar coordinates, via the transformation $x = p(r, \theta) = r \cos \theta$, $y = q(r, \theta) = r \sin \theta$. Then the Jacobian is given by $\det \begin{pmatrix} \partial r \cos \theta / \partial r & \partial r \cos \theta / \partial \theta \\ \partial r \sin \theta / \partial r & \partial r \sin \theta / \partial \theta \end{pmatrix} = \det \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} = r$. Now, Change of Variables gives $I^2 = \int_0^{\tau} \int_0^{\infty} e^{-r^2(\cos^2 \theta + \sin^2 \theta)} r dr d\theta = \int_0^{\tau} [-\frac{1}{2} e^{-r^2}]_0^{\infty} d\theta = \int_0^{\tau} \frac{1}{2} d\theta = \tau/2$.

This theorem, whose use is second nature to applied mathematicians and probability theorists, was surprisingly resistant to formal proof. Victor Katz attributes its first completely satisfactory treatment to Elie Cartan in the 1890s, over 125 years after Leonhard Euler first proposed the technique while inventing double integration.

Web link: mathinsight.org/double_integral_change_variables_introduction

Further reading: *The Genius of Euler* by William Dunham (ed.), the Mathematical Association of America, 2007 (Katz's authoritative paper is reproduced in part 2). The Kelvin quote is from Silvanus P. Thompson's *Life of Lord Kelvin, vol.2*, OUP.

