




# THEOREM OF THE DAY

**The Lagrange Interpolation Formula** Given  $n$  distinct real values,  $x_1, \dots, x_n$ ,  $n \geq 2$ , and any  $n$  points,  $(x_1, y_1), \dots, (x_n, y_n)$ , in the Cartesian plane, there is unique polynomial curve,  $y = p(x)$ , of degree  $n - 1$ , passing these points, specified by

$$p(x) = \sum_{i=1}^n y_i \prod_{i \neq j} (x - x_j) / (x_i - x_j).$$



## Remarks:

1. Writing the formula explicitly for  $n = 2$  points gives  $y = y_1(x - x_2)(x_1 - x_2)^{-1} + y_2(x - x_1)(x_2 - x_1)^{-1}$ , the equation for the unique straight line passing through these points: with a little manipulation, it becomes the more memorably symmetrical straight line equation  $(y - y_1)/(y_2 - y_1) = (x - x_1)/(x_2 - x_1)$ .
2. The  $n = 2$  calculation looks very similar to that which solves the Chinese Remainder Theorem for two modular equations, and indeed there is a close connection.
3. The calculation also reveals why the formula works in general: each term is a polynomial which takes the value  $y_i$  when  $x = x_i$  and is zero when  $x = x_j$ ,  $j \neq i$ .
4. We can deduce uniqueness thus: suppose  $p(x)$  and  $q(x)$  are polynomials of degree  $n - 1$  through our  $n$  points, and define the polynomial  $r(x) = p(x) - q(x)$ . Now for  $i = 1, \dots, n$ ,  $r(x_i) = 0$  so, by the Factor Theorem,  $(x - x_i)$  is a factor of  $r(x)$ . But this means  $r(x)$  has a factor  $(x - x_1)(x - x_2) \cdots (x - x_n)$  and must therefore be the zero polynomial, otherwise it has higher degree than  $p$  and  $q$  which is impossible.

Edward Waring extracted this formula in 1779 from a more general one of Newton. Independently, Lagrange did likewise in 1795.

**Web link:** [imagescience.org/meijering/publications/1015/](http://imagescience.org/meijering/publications/1015/)

**Further reading:** *Over and Over Again* by Gengzhe Chang and Thomas W. Sederberg, MAA, 1998, chapter 17.

