







# THEOREM OF THE DAY

**Brahmagupta's Formula** *The area  $K$  of a cyclic quadrilateral with side lengths  $a, b, c, d$  and semiperimeter  $s = (a + b + c + d)/2$  is given by*

$$K = \sqrt{(s - a)(s - b)(s - c)(s - d)}.$$



(i) Euclid of Alexandria  
*Elements, Book IV, (c. 300 BC)*  
Circumcircling the triangle

(ii) Heron of Alexandria  
*Metrica, (1st century AD)*  
Sides  $\rightarrow$  area (triangle)

(iii) Brahmagupta  
*The Brāhmasphutasiddhānta, (628 AD)*  
Sides  $\rightarrow$  area (cyclic quadrilateral)

(iv) J.L. Coolidge  
*Amer. Math. Monthly, (1939)*  
Sides  $\rightarrow$  area (any quadrilateral)

(i) The familiar formula for triangular area ( $1/2 \times \text{base} \times \text{height}$ ) was known to Greek mathematicians for at least three hundred years before Euclid catalogued all of known geometry in his *Elements*, including (Book IV, Proposition 5) the construction of the unique *circumcircle* which passes through the vertices of a triangle, by intersecting the perpendicular bisectors of its sides.

(ii) Three hundred years after *that* came the famous **Heron's formula**: *the area  $K$  of a triangle with sides  $A, B, C$  and semiperimeter  $s = (A + B + C)/2$  is given by  $K = \sqrt{(s - A)(s - B)(s - C)s}$ .*

(iii) Although Greek mathematics was apparently unknown to medieval Indian (as opposed to Islamic) scholars, Brahmagupta effectively put Heron's triangle back into the circle: take two non-overlapping circumscribed triangles sharing a common edge ( $C$  in the picture); the result is a *cyclic quadrilateral*, one whose vertices all lie on a circle. And now the area of the quadrilateral replaces the final  $s$  in Heron's formula by  $s - d$ . If point  $P$  is allowed to approach point  $Q$  then  $d$  becomes zero and  $c$  becomes  $C$ , recovering Heron.

(iv) Another thirteen hundred years pass and the circumscribing circle is removed once more in American mathematician Julian Lowell Coolidge's **quadrilateral area formula**: *the area of an arbitrary convex quadrilateral with sides  $a, b, c, d$ ,  $a$  opposite to  $d$ , with diagonals  $e, f$ , and with semiperimeter  $s$ , is given by  $K = \sqrt{(s - a)(s - b)(s - c)(s - d) - \frac{1}{4}(ad + bc + ef)(ad + bc - ef)}$ . This generalises Brahmagupta by virtue of another classic of antiquity, **Ptolemy's Theorem**: *quadrilateral  $a, b, c, d$ ,  $a$  opposite to  $d$ , with diagonals  $e, f$ , is cyclic if and only if  $ad + bc = ef$ .**

Brahmagupta's formula appears in his *Brāhmasphutasiddhānta*, a treatise on astronomy. Brahmagupta's writings contain the first known treatment of zero and negative numbers.

**Web link:** [www.mathpages.com/home/kmath196/kmath196.htm](http://www.mathpages.com/home/kmath196/kmath196.htm)

**Further reading:** *Journey Through Genius* by William Dunham, John Wiley & Sons, 1990.

Created by Robin Whitty for [www.theoremoftheday.org](http://www.theoremoftheday.org)