

# The Polygonal Number Theorem

Robin Whitty


[theoremoftheday.org](http://theoremoftheday.org)

6 May, 2021

# Polygonal Numbers

For  $m \geq 1$ , the  $k$ -th polygonal number of order  $m + 2$  is defined to be

$$P_m(k) = \frac{m}{2} (k^2 - k) + k, k \geq 0.$$



From  $m + 1$  vertices of an  $(m + 2)$ -gon, add new vertices along 'rays', interpolating  $1, 2, 3, \dots$ , vertices.

For  $m = 3$ , pentagonal numbers:  $1, 5, 12, 22, \dots$

# Triangular Numbers

For  $m = 1$ , the polygonal numbers of order 3 are the triangular numbers

$$P_1(k) = \frac{1}{2}(k^2 + k), k \geq 0.$$

1										
1	1									
1	2	1								
1	3	3	1							
1	4	6	4	1						
1	5	10	10	5	1					
1	6	15	20	15	6	1				
1	7	21	35	35	21	7	1			
1	8	28	56	70	56	28	8	1		
1	9	36	84	126	126	84	36	9	1	
1	10	45	120	210	252	210	120	45	10	1
			...							

The triangular numbers appear on the 2nd diagonal of Pascal's triangle. (The sum of their reciprocals converges, as is the case with all diagonals except the 0th and 1st - Nick Hobson, M500, Issue 216.)


# Fermat's Polygonal Number Conjecture

Fermat, in a letter to Mersenne in 1636, asserted that, for all positive integer  $m$ , every nonnegative integer is a sum of  $m + 2$  polygonal numbers of order  $m + 2$

$$23 = \begin{array}{c} \circ \\ \circ \circ \\ \circ \circ \circ \end{array} + \begin{array}{c} \circ \\ \circ \circ \\ \circ \circ \circ \\ \circ \circ \circ \circ \end{array} + \begin{array}{c} \circ \\ \circ \circ \\ \circ \circ \circ \\ \circ \circ \circ \circ \\ \circ \circ \circ \circ \circ \end{array}$$

# Fermat $m = 1$ : Gauss's Eureka Theorem


Gauss proved the case  $m = 1$  of Fermat's conjecture on July 10, 1796, as per the following entry in his diary:



Equivalently, if  $n \equiv 3 \pmod{8}$  then  $n$  can be written as  $n = x^2 + y^2 + z^2$  for odd integers  $x, y, z$ .

# Tetrahedral Numbers


The third diagonal of Pascal's triangle are the **tetrahedral numbers**, being the number of spheres that can be densely packed in a triangular pyramid.



1											
1	1										
1	2	1									
1	3	3	1								
1	4	6	4	1							
1	5	10	10	5	1						
1	6	15	20	15	6	1					
1	7	21	35	35	21	7	1				
1	8	28	56	70	56	28	8	1			
1	9	36	84	126	126	84	36	9	1		
1	10	45	120	210	252	210	120	45	10	1	
			...								

The diagonals generally generalise the triangular numbers to **figurate** numbers: tetrahedral, 4-simplex, etc, although definitions vary.


# Figurate Numbers



Notations	ix
Preface	xv
1. Plane figurate numbers	1
1.1 Definitions and formulas . . . . .	1
1.2 Main properties of polygonal numbers . . . . .	12
1.3 Square triangular numbers . . . . .	22
1.4 Other highly polygonal numbers . . . . .	35
1.5 Amount of a given number in all polygonal numbers . . . . .	45
1.6 Centered polygonal numbers . . . . .	47
1.7 Other plane figurate numbers . . . . .	61
1.8 Generalized plane figurate numbers . . . . .	76
2. Space figurate numbers	87
2.1 Pyramidal numbers . . . . .	88
2.2 Cubic numbers . . . . .	99
2.3 Octahedral numbers . . . . .	105
2.4 Other regular polyhedral numbers . . . . .	109
2.5 Some semiregular and star polyhedral numbers . . . . .	115
2.6 Centered space figurate numbers . . . . .	120

## Fermat $m = 2$ : Square Numbers

When  $m = 2$  we have the polygonal numbers of order 4, the **square numbers**. Any positive integer may be written as a sum of at most four square numbers. This was known Diophantus of Alexandria and was first explicitly asserted by Claude Gaspard Bachet de Méziriac, who translated Diophantus's *Arithmetica* into Latin in 1621. The first proof is due to Lagrange in 1770.



Any collection of blocks may be arranged as a square pyramid of height at most 4 blocks. E.g.  $23 = 3^2 + 3^2 + 2^2 + 1^2$ .



# The Fifteen Theorem

In 1993, John Conway and his student William Schneeberger proved:

*If a positive-definite quadratic form defined by a symmetric, integral matrix takes each of the values 1, 2, 3, 5, 6, 7, 10, 14, 15, then it takes all positive integer values.*

	$w$	$x$	$y$	$z$
$w$	1			
$x$		1		
$y$			1	
$z$				1

$$w^2 + x^2 + y^2 + z^2$$

$w$	$x$	$y$	$z$	
0	0	0	1	1
0	0	1	1	2
0	1	1	1	3
0	0	1	2	5
0	1	1	2	6
1	1	1	2	7
1	1	2	2	10
0	1	2	3	14
1	1	2	3	15

## Waring's Problem

Coincidentally with Lagrange's proof of the four-squares theorem, Edward Waring proposed (and, in 1909, Hilbert confirmed, nonconstructively):

*For each positive integer  $k$ , there is a positive integer  $g(k)$  such that every nonnegative integer may be written as a sum of at most  $s$  integers raised to power  $k$ .*

E.g.  $23 = 2^3 + 2^3 + 1^3 + 1^3 + 1^3 + 1^3 + 1^3 + 1^3 + 1^3$ : one of only two cases where  $g(3) = 9$  nontrivial cubes are required. It is thought that six are sufficient for large  $n$  ( $G(3) = 6$ ).

**Theorem (1940s):** If  $\left\{ \left( \frac{3}{2} \right)^n \right\} \leq 1 - \left( \frac{3}{4} \right)^n$ , where  $\{ \cdot \}$  denotes the fractional part of a real number, then

$$g(n) = 2^n + \left[ \left( \frac{3}{2} \right)^n \right] - 2,$$

where  $[ \cdot ]$  denotes the integer part of a real number.

## Fermat $m \geq 3$ : Cauchy's Lemma

Cauchy (1815) showed that Gauss's Eureka Theorem implies:

*If  $a$  and  $b$  are odd positive integers satisfying*

$$b^2 - 4a < 0 \quad \text{and} \quad 0 < b^2 + 2b - 3a + 4,$$

*then there exist nonnegative integers  $s, t, u, v$  such that*

$$a = s^2 + t^2 + u^2 + v^2 \quad \text{and} \quad b = s + t + u + v.$$

E.g.  $a = 259, b = 29,$

$$b^2 - 4a = 841 - 1036 < 0$$

$$b^2 + 2b - 3a + 4 = 841 + 58 - 777 + 4 > 0$$

$$259 = 13^2 + 7^2 + 5^2 + 4^2$$

$$29 = 13 + 7 + 5 + 4$$

## Fermat $m \geq 3$ : Cauchy's Theorem

Cauchy (1815): If  $m \geq 3$  then every nonnegative number can be written as a sum of at most  $m + 2$  polygonal numbers of order  $m + 2$ .

**E.g.**

$$375 = P_3(13) + P_3(7) + P_3(5) + P_3(4) + P_3(1) = 247 + 70 + 35 + 22 + 1.$$

(Recall

$$P_m(k) = \frac{m}{2} (k^2 - k) + k, k \geq 0.)$$

Subsequent work, notably by Jean François Théophile Pépin, constructed explicit order  $m + 2$  polygonal representations for all integers  $n < 120m$ .

## Fermat $m \geq 3$ : Melvyn Nathanson's proof

1. Assume that  $m \geq 3$ . Choose an odd positive integer  $b$  such that
  - 1.1 We can write  $n \equiv b + r \pmod{m}$ ,  $0 \leq r \leq m - 2$ ; and
  - 1.2 If  $a = 2 \left( \frac{n - b - r}{m} \right) + b$ , an odd positive integer by virtue of (1.1), then

$$b^2 - 4a < 0 \quad \text{and} \quad 0 < b^2 + 2b - 3a + 4. \quad (*)$$

2. Invoke **Cauchy's Lemma**: *If  $a$  and  $b$  are odd positive integers satisfying  $(*)$  then there exist nonnegative integers  $s, t, u, v$  such that*

$$a = s^2 + t^2 + u^2 + v^2 \quad \text{and} \quad b = s + t + u + v.$$

3. From the definition of  $a$  in (1.2), write  $n = \frac{m}{2}(a - b) + b + r$   
 $= \frac{m}{2}(s^2 - s) + s + \dots + \frac{m}{2}(v^2 - v) + v + r.$

## Fermat $m \geq 3$ : Nathanson's proof, the small print

1. Assume that  $m \geq 3$ . Choose an odd positive integer  $b$  such that

1.1 We can write  $n \equiv b + r \pmod{m}$ ,  $0 \leq r \leq m - 2$ ; and


1.2 If  $a = 2 \left( \frac{n - b - r}{m} \right) + b$ , an odd positive integer by virtue of (1.1), then

$$b^2 - 4a < 0 \quad \text{and} \quad 0 < b^2 + 2b - 3a + 4. \quad (*)$$

For a given  $n$  and  $m$  an interval for  $b$  may be expressed, via the quadratic formula, purely in terms of  $n$  and  $m$ . Namely, the interval  $[1/2 + \sqrt{6(n/m) - 3}, 2/3 + \sqrt{8(n/m) - 8}]$  is guaranteed to be bounded by the zeros of the quadratics and to have length at least 4 for  $n \geq 120m$ .

Any interval of length 4 must contain two odd integers which together contain a complete set of residues mod  $m$ .

# Nathanson's Interval



## Legendre's Theorem (1832)

As  $n$  gets larger, the gap between the roots of the two quadratics in Nathanson's proof, as illustrated in the preceding slide, gets larger. This means the congruence  $b \equiv b + r$  can be achieved with smaller and smaller values of  $r$ . Eventually  $r = 0$  is always possible, for odd  $m$ , while for even  $m$  the least possible  $r$  oscillates between 0 and 1. Since  $r$  contributes all the polygonal numbers in Nathanson's proof except the first four, this gives an improvement on Cauchy's 1815 theorem, due to Legendre:

*Let  $m > 3$ . If  $m$  is odd, then every sufficiently large integer is the sum of four polygonal numbers of order  $m + 2$ . If  $m$  is even, then every sufficiently large integer is the sum of five polygonal numbers of order  $m + 2$ , one of which is either 0 or 1.*



## Some links

- ▶ Nick Hobson, "Solution 213.1 - Pascal triangle sums", M500, Issue 216, pp. 1–2, [m500.org.uk/magazine/](http://m500.org.uk/magazine/).
- ▶ Lagrange's Four-Squares Theorem at [theoremoftheday.org](http://theoremoftheday.org):  
[www.theoremoftheday.org/Theorems.html#11](http://www.theoremoftheday.org/Theorems.html#11)
- ▶ Elena Deza and Michel Marie Deza, *Figurate Numbers*, World Scientific, 2012. Details and online reviews at [www.theoremoftheday.org/Resources/Bibliography.htm#ElenaDeza](http://www.theoremoftheday.org/Resources/Bibliography.htm#ElenaDeza)
- ▶ The Fifteen Theorem at [theoremoftheday.org](http://theoremoftheday.org):  
[www.theoremoftheday.org/Theorems.html#79](http://www.theoremoftheday.org/Theorems.html#79)
- ▶ The Polygonal Number Theorem and Nathanson's proof are described here at [theoremoftheday.org](http://theoremoftheday.org):  
[www.theoremoftheday.org/Theorems.html#262](http://www.theoremoftheday.org/Theorems.html#262)